CONTENTS

Preface vi
Committees viii
Variance analysis fifteen years ago and now
A. Péneloux, E. Neau and A. Gramajo (Marseille, France) 1–16
Combined excess free energy models and equations of state
R.A. Heidemann and S.L. Kokal (Calgary, Alb., Canada) 17–37
A non-cubic equation of state for *PVT* and phase equilibrium calculations (pure compounds)
G. Soave (San Donato Milanese, Italy) 39–57
A group contribution method for prediction of hydrocarbon saturated liquid volumes
M. Rogalski and E. Neau (Marseille, France) 59–69
A modified volume-translated Peng–Robinson equation with temperature dependent parameters
W. Sheng and B.C.-Y Lu (Ottawa, Ont., Canada) 71–80
Generalization of binary interaction parameters of Peng–Robinson equation of state for systems containing hydrogen
H. Nishiumi and H. Gotoh (Tokyo, Japan) 81–88
Prediction of phase equilibria for hydrocarbon systems in a wide range of pressure and temperature by AEOS equation of state
B. Wiśniewska and S. Malanowski (Warsaw, Poland) 89–95
Capabilities of the PFGC–MES EOS in predicting the VLE data of refrigerants and light hydrocarbon mixtures
M.H. Shariat, N. Mehranbod and M. Moshfeghian (Shiraz, Iran) 97–104
VLE correlation for binary 1-alkanol + n-alkane mixtures
M.R. Margerum and B.C.-Y Lu (Ottawa, Ont., Canada) 105–118
Thermophysical properties of n-alkanes from C1 to C20 and their prediction for higher ones
K. Magoulas and D. Tassios (Athens, Greece) 119–140
Correlation and prediction of VLE data for n-alkane mixtures
R. Stryjek (Warsaw, Poland) 141–152
Correlation of the critical properties of alkanes and alkanols
A.S. Teja, R.J. Lee, D. Rosenthal and M. Anselme (Atlanta, GA, U.S.A.) . 153–169
Thermodynamics of associated solutions. Mixtures of 1-alkanols
C.G. Panayiotou (Thessaloniki, Greece) 171–188
VLE of alkanol–alkane systems with the modified CRG model
E. Neau, M. Rogalski (Marseille, France), P. Alessi and M. Fermeglia (Trieste, Italy) 189–202
Correlation of the spectroscopic and thermodynamic properties of solutions of propanols or butanols with non-associating components using the UNIQUAC associated-solution model
I. Nagata and K. Miyamoto (Ishikawa, Japan) 203–218

Analysis of excess enthalpies of ethyl formate + n-alkane or 1-alkanol with two group contribution models
J. Ortega (Las Palmas, Spain), J.L. Legido, J. Fernández, M. López, L. Pías and M.I. Paz Andrade (Santiago de Compostela, Spain) 219–234
Simultaneous correlation of vapor–liquid equilibrium and heat of mixing data for ternary systems
G. Figurski (Halle, G.D.R.), J. Gregorowicz and S. Malanowski (Warsaw, Poland) 235–247
Infinite dilution activity coefficients in poly(ethylene glycol)
E.L. Sørensen, W. Hao (Lyngby, Denmark) and P. Alessi (Trieste, Italy) . 249–256
Prediction of phase equilibria in polymer systems
G. Carvoli, A. Castelli (Novara, Italy) and A.M. Marconi (Mantua, Italy) .. 257–267
Thermodynamic consistency test of vapor–liquid equilibrium data – methanol–water, benzene–cyclohexane and ethyl methyl ketone–water
K. Kojima, H.M. Moon and K. Ochi (Tokyo, Japan) 269–284
A data base standard for the evaluation of vapor–liquid-equilibrium models
R.P. Danner and M.A. Gess (University Park, PA, U.S.A.) 285–301
Isothermal vapour–liquid equilibrium data for the system n-pentane/n-propanol
P. Rice (Loughborough, Gt. Britain), A. El-Nikheli (Riyadh, Saudi Arabia) and A.S. Teja (Atlanta, GA, U.S.A.) 303–312
Solubility of H_2S, CO_2, CH_4 and C_2H_6 in sulfolane at elevated pressures
F.-Y. Jou, R.D. Deshmukh, F.D. Otto and A.E. Mather (Edmonton, Alb., Canada) 313–324
Binary and ternary phase behaviour of the system water–ethanol–carbon dioxide
E. Martinez de la Ossa (Cadiz, Spain), V. Brandani, G. Del Re, G. Di Giacomo and E. Ferri (Aquila, Italy) 325–340
Measurement and correlation of liquid–liquid equilibria up to critical solution temperature
K. Ochi, M. Tada and K. Kojima (Tokyo, Japan) 341–359
Author Index 361